

IFT-P.002/2000

FOUR-POINT AMPLITUDE FROM OPEN SUPERSTRING FIELD THEORY

Nathan Berkovits¹
and
Carlos Tello Echevarria ²

*Instituto de Física Teórica, Universidade Estadual Paulista
Rua Pamplona 145, 01405-900, São Paulo, SP, Brasil*

December 1999

Abstract

An open superstring field theory action has been proposed which does not suffer from contact term divergences. In this paper, we compute the on-shell four-point tree amplitude from this action using the Giddings map. After including contributions from the quartic term in the action, the resulting amplitude agrees with the first-quantized prescription.

¹e-mail: nberkovi@ift.unesp.br

²e-mail: carlos@ift.unesp.br

1 Introduction

To study non-perturbative features of superstring theory, a field theory action for the superstring could be useful. As shown by Wendt in 1989 [1], Witten's cubic action [2] for open superstring field theory has divergent contact term problems due to colliding interaction-point operators. If the Neveu-Schwarz string field A carries picture -1 , the cubic action is $\int(AQA + ZA^3)$ where $Z = \{Q, \xi\}$ is the picture-raising operator which is inserted at the interaction-point [2]. Since Z has a singularity with itself, the four-point amplitude computed using the above action has contact term divergences which break gauge invariance and make the action inconsistent. Similar contact term problems exist with other choices of picture [3] and with other open superstring field theory actions such as the light-cone and covariantized light-cone actions.[4]

In 1995, an action for open superstring field theory was proposed which does not suffer from contact term divergences [5]. This action was constructed by embedding the $N=1$ superstring into the $N=2$ superstring and resembles a Wess-Zumino-Witten action. Using modified Green-Schwarz variables, this open superstring field theory action can be written in a manifestly $SO(3,1)$ super-Poincaré invariant manner [5]. This spacetime-supersymmetric version of the action is easily generalized to any compactification to four dimensions which preserves at least $N=1$ $d = 4$ supersymmetry. It is also possible to write this field theory action using Ramond-Neveu-Schwarz variables and, at least in the Neveu-Schwarz sector, this can be done in a manifestly $SO(9,1)$ Lorentz invariant manner.

In this paper, we shall use this action to explicitly compute the four-point open superstring tree amplitude. When the external states are on-shell, this amplitude will be shown to agree with the first-quantized result. There are no contact term divergences, and the quartic vertex plays a crucial role in cancelling a *finite* contact term contribution coming from two cubic vertices.

In section 2, we shall review the Wess-Zumino-Witten-like action for open superstring field theory. In section 3, we shall discuss gauge fixing and the Giddings map for computing four-point amplitudes using open string field theory. In section 4, we shall explicitly compute the four-point on-shell tree amplitude and will prove equivalence with the first-quantized result.

2 Review of Superstring Field Theory Action

For any critical ($\hat{c} = 2$) N=2 superconformal representation with N=2 generators $[T, G^+, G^-, J]$, one can construct the following open string field theory action:

$$S = \frac{1}{2} \int [(e^{-\Phi} G_0^+ e^{\Phi})(e^{-\Phi} \tilde{G}_0^+ e^{\Phi}) - \int_0^1 dt (e^{-t\Phi} \partial_t e^{t\Phi}) \{e^{-t\Phi} G_0^+ e^{t\Phi}, e^{-t\Phi} \tilde{G}_0^+ e^{t\Phi}\}]. \quad (1)$$

which resembles the Wess-Zumino-Witten action $\int [(g^{-1} dg)^2 + \int dt (g^{-1} dg)^3]$. The open string fields Φ are glued together using Witten's midpoint interaction [6] and are restricted to be U(1)-neutral with respect to J . The open string fields carry Chan-Paton factors which will be suppressed throughout this paper.

The fermionic operators G_0^+ and $\tilde{G}_0^+$ are constructed from the N=2 superconformal generators in the following manner [7]: After twisting, the fermionic generator G^+ carries conformal weight +1 and the fermionic generator G^- carries conformal weight +2. Furthermore, after bosonizing the U(1) generator $J = \partial H$, one can construct SU(2) currents $[J^{++} = e^{iH}, J = \partial H, J^{--} = e^{-iH}]$ of conformal weight $[0, 1, 2]$ after twisting. Commuting the zero modes of these SU(2) currents with the fermionic generators, one obtains two new fermionic generators, $\tilde{G}^+ = [G^-, J_0^{++}]$ and $\tilde{G}^- = [G^+, J_0^{--}]$, of conformal weight +1 and +2 respectively. G_0^+ is defined to be the zero mode of G^+ and $\tilde{G}_0^+$ is defined to be the zero mode of $\tilde{G}^+$. Note that $\{G_0^+, \tilde{G}_0^+\} = 0$ and after twisting, the U(1) anomaly implies that non-vanishing correlation functions on a sphere must carry +2 U(1) charge.

The action of (1) can be justified by considering the following three examples of critical N=2 superconformal representations. The first example is the open self-dual string where Φ depends on the string modes of the variables $[x^{\alpha\dot{\alpha}}, \psi^\alpha, \bar{\psi}^{\dot{\alpha}}]$ for $\alpha, \dot{\alpha} = 1$ to 2. For this N=2 superconformal representation, $G^+ = \psi_\alpha \partial x^{\alpha+}$, $\tilde{G}^+ = \psi_\alpha \partial x^{\alpha-}$, and $J = \psi^\alpha \bar{\psi}_\alpha$. Since there are no massive physical fields, the dependence of Φ on the non-zero modes of $[x^{\alpha\dot{\alpha}}, \psi^\alpha, \bar{\psi}^{\dot{\alpha}}]$ gives rise to auxiliary and gauge fields. The massless physical field comes from dependence on the zero mode of $x^{\alpha\dot{\alpha}}$, $\Phi(x_0^{\alpha\dot{\alpha}})$, which when plugged into (1) reproduces the Donaldson-Nair-Schiff action [8] for D=4 self-dual Yang-Mills.

The second example is the N=2 embedding of the superstring using the standard RNS worldsheet variables $[x^\mu, \psi^\mu, b, c, \xi, \eta, \phi]$ where $\mu = 0$ to 9 and

the super-reparameterization ghosts have been fermionized as $\beta = e^{-\phi}\partial\xi$ and $\gamma = e^{\phi}\eta$ [7]. In this example, the Neveu-Schwarz contribution to the field theory action is obtained by defining $\Phi = \xi_0 A$ where A is the standard Neveu-Schwarz string field in the -1 picture. (The Ramond contribution to the superstring field theory action is not yet known.) In the N=2 embedding of the superstring, G_0^+ is the BRST charge Q and $\tilde{G}_0^+$ is the zero mode of η . So the linearized equation of motion coming from the action of (1) is

$$0 = G_0^+ \tilde{G}_0^+ \Phi = Q\eta_0(\xi_0 A) = QA$$

as desired. Furthermore, since $J = bc + \xi\eta$, the U(1)-neutrality condition implies that Φ has zero RNS ghost-number.

The third example is the ‘hybrid’ description of the superstring using the four-dimensional superspace variables $[x^m, \theta^\alpha, \bar{\theta}^{\dot{\alpha}}]$ where $m=0$ to 3 and $\alpha, \dot{\alpha} = 1$ to 2, combined with a chiral boson ρ and the variables of a $c = 9$ N=2 superconformal field theory which describes a six-dimensional compactification manifold [9]. In this example, the compactification-independent part of the field theory action is obtained by defining Φ to be a string field depending only on the four-dimensional superspace variables. (The compactification-dependent contribution to the action was worked out in [5] and involves two other string fields in addition to Φ .) The massless and first massive level of this action have been explicitly computed in [5] and [10] and describe a super-Yang-Mills and massive spin-2 multiplet in N=1 $d = 4$ superspace.

3 Gauge Fixing and the Giddings Map

Similar to the Wess-Zumino-Witten action, the action of (1) is invariant under the gauge transformation

$$\delta e^\Phi = (G_0^+ \Omega) e^\Phi + e^\Phi (\tilde{G}_0^+ \tilde{\Omega})$$

where Ω and $\tilde{\Omega}$ are arbitrary string fields of -1 U(1)-charge. At linearized level, this transformation reduces to $\delta\Phi = G_0^+ \Omega + \tilde{G}_0^+ \tilde{\Omega}$ which allows the gauge-fixing conditions

$$G_0^- \Phi = \tilde{G}_0^- \Phi = 0 \tag{2}$$

where G_0^- and $\tilde{G}_0^-$ are the zero modes of G^- and $\tilde{G}^- = [G^+, J_0^-]$. The gauge-fixing conditions of (2) can be obtained, for example, by choosing

$$\Omega = -(T_0)^{-1} G_0^- \Phi, \quad \tilde{\Omega} = -(T_0)^{-2} \tilde{G}_0^- G_0^+ \Phi,$$

so that $G_0^- \delta \Phi = -G_0^- \Phi$ and $\tilde{G}_0^- \delta \Phi = -\tilde{G}_0^- \Phi$ where we have used that $\{G_0^-, G_0^+\} = \{\tilde{G}_0^-, \tilde{G}_0^+\} = T_0$.

In this gauge, the linearized propagator is $\mathcal{P} = (T_0)^{-2} G_0^- \tilde{G}_0^-$ since $\tilde{G}_0^+ G_0^+ \Phi = 0$ is the linearized equation of motion and $\mathcal{P} \tilde{G}_0^+ G_0^+ \Phi = \Phi$. Although this propagator looks complicated, we shall show in section 4 that it can be simplified to $\mathcal{P} = (T_0)^{-1} J_0^{--}$ when computing on-shell tree amplitudes.

Although the action of (1) contains vertices with arbitrary numbers of string fields, only the cubic and quartic vertices will be necessary for computing four-point tree amplitudes. Expanding the action of (1), one obtains

$$S = \int \left(\frac{1}{2} G_0^+ \Phi \tilde{G}_0^+ \Phi - \frac{1}{6} \Phi \{G_0^+ \Phi, \tilde{G}_0^+ \Phi\} - \frac{1}{24} [\Phi, G_0^+ \Phi] [\Phi, \tilde{G}_0^+ \Phi] + \dots \right). \quad (3)$$

Using Witten's gluing prescription [6] for string fields, the cubic and quartic vertex from (3) are described by the diagrams




Figure 1

For four-point tree amplitudes, contributions can come from two cubic vertices or from one quartic vertex. The Witten diagram for two cubic vertices connected by a propagator of length τ is given by




Figure 2

where the propagator is integrated along the contour c . The Witten diagram for a quartic vertex is given by Figure 2 in the limit that $\tau \rightarrow 0$.

As was shown by Giddings [11], it is convenient to perform a Schwarz-Christoffel transformation from the diagram of figure 2 to the upper half plane such that the four external strings are mapped to the points $\pm\alpha(\tau)$ and $\pm(\alpha(\tau))^{-1}$. If strings $[(1), (2), (3), (4)]$ are mapped to the points $[-\alpha, \alpha, \alpha^{-1}, -\alpha^{-1}]$ in this order, then $0 < \alpha(\tau) \leq \delta$ where $\delta \equiv \sqrt{2} - 1$, $\alpha(\tau = 0) = \delta$ and $\alpha(\tau = \infty) = 0$. On the other hand, if strings $[(1), (2), (3), (4)]$ are mapped to the points $[-\alpha^{-1}, -\alpha, \alpha, \alpha^{-1}]$ in this order, then $\delta \leq \alpha(\tau) < 1$ where $\alpha(\tau = 0) = \delta$ and $\alpha(\tau = \infty) = 1$.

4 Computation of Four-Point Amplitude

The tree-level scattering amplitude for four external string fields labeled by the letters $[A, B, C, D]$ gets contributions either from the diagram of Figure 2 or from the second diagram of Figure 1. There are 24 different ways to match the string fields $[A, B, C, D]$ with the external legs $[(1), (2), (3), (4)]$ and we shall restrict our attention to those 4 combinations where $[A, B, C, D]$ are cyclically ordered. For contributions coming from the diagram of Figure 2, these 4 combinations split into two ‘s-channel’ contributions, $\mathcal{A}_s$, where string A is associated with leg (2) or leg (4) and two ‘t-channel’ contributions, $\mathcal{A}_t$, where string A is associated with leg (1) or leg (3). The four cyclically related combinations coming from the quartic vertex of Figure 1 will be called $\mathcal{A}_q$.

In this section, it will be shown that the sum of these three contributions, $\mathcal{A} = \mathcal{A}_s + \mathcal{A}_t + \mathcal{A}_q$, reproduces the first-quantized result for the on-shell scattering amplitude which can be written as [7]

$$\mathcal{A} = \int_0^1 d\alpha \langle \left(\int d^2 z \mu_\alpha(z, \bar{z}) G^-(z) \right) \quad (4)$$

$$\Phi_A(-\alpha^{-1}) G_0^+ \Phi_B(-\alpha) G_0^+ \Phi_C(\alpha) \tilde{G}_0^+ \Phi_D(\alpha^{-1}) \rangle.$$

where $\mu_\alpha(z, \bar{z})$ is the appropriate Beltrami differential for an α -dependent parameterization of the modulus and $\langle \rangle$ signifies the two-dimensional correlation function in the upper half-plane. To relate (4) to the standard expression for the scattering amplitude of four Neveu-Schwarz strings, recall from section 2 that $\Phi = \xi_0 V^{(-1)}$ where $V^{(-1)}$ is the Neveu-Schwarz vertex operator in the -1 picture. Furthermore, $G^- = b$, $\tilde{G}_0^+ \Phi = \eta_0 \xi_0 V^{(-1)} = V^{(-1)}$ and

$G_0^+ \Phi = Q \xi_0 V^{(-1)} = V^{(0)}$ where $V^{(0)}$ is the Neveu-Schwarz vertex operator in the zero picture. So (4) implies that

$$\mathcal{A} = \int_0^1 d\alpha \langle \left(\int d^2 z \mu_\alpha(z, \bar{z}) b(z) \right) \xi_0 V_A^{(-1)}(-\alpha^{-1}) V_B^{(0)}(-\alpha) V_C^{(0)}(\alpha) V_D^{(-1)}(\alpha^{-1}) \rangle,$$

which agrees with the first-quantized prescription of [12] in the large Hilbert space, i.e. in the Hilbert space including the ξ zero mode.

4.1 Computation of s-channel contribution

We shall first compute the contribution $\mathcal{A}_s$ from s-channel diagrams. Using the cubic vertices and propagator $\mathcal{P}$ from section 3 and including the various combinatorial factors,

$$\mathcal{A}_s = 2(3)^2 \frac{1}{2} \left(-\frac{1}{6}\right)^2 \langle (G_0^+ \Phi_A(4) \tilde{G}_0^+ \Phi_B(1) + \tilde{G}_0^+ \Phi_A(4) G_0^+ \Phi_B(1))$$

$$\mathcal{P} (G_0^+ \Phi_C(2) \tilde{G}_0^+ \Phi_D(3) + \tilde{G}_0^+ \Phi_C(2) G_0^+ \Phi_D(3)) \rangle_W$$

where $\langle \rangle_W$ signifies the two-dimensional correlation function in the Witten diagram of Figure 2. Note that one can always choose the G_0^+ and $\tilde{G}_0^+$ operators of the cubic vertices to act on external legs since

$$\begin{aligned} & \langle (G_0^+ \Phi_A(1) \tilde{G}_0^+ \Phi_B(2) + \tilde{G}_0^+ \Phi_A(1) G_0^+ \Phi_B(2)) \Phi_C(3) \rangle = \\ & \langle \Phi_A(1) (\tilde{G}_0^+ \Phi_B(2) G_0^+ \Phi_C(3) + G_0^+ \Phi_B(2) \tilde{G}_0^+ \Phi_C(3)) \rangle \end{aligned}$$

by deforming the contour integral of G^+ and $\tilde{G}^+$ off of $\Phi_A(1)$.

The propagator $\mathcal{P} = (T_0)^{-2} G_0^- \tilde{G}_0^-$ can be simplified by writing $\tilde{G}_0^- = [G_0^+, J_0^{--}]$ and deforming the contour integral of G^+ off of J_0^{--} . When the external string fields are on-shell, i.e. $G_0^+ \tilde{G}_0^+ \Phi = 0$, the contour integral of G^+ only contributes by hitting the remaining G_0^- of the propagator. Since $\{G_0^+, G_0^-\} = T_0$, this means that the propagator can be simplified to $\mathcal{P} = (T_0)^{-1} J_0^{--}$ when all external string fields are on-shell. As usual, it is convenient to rewrite $(T_0)^{-1} = \int_0^\infty d\tau e^{-\tau T_0}$ so the s-channel contribution is

$$\begin{aligned} \mathcal{A}_s = \frac{1}{4} \int_0^\infty d\tau \langle \int_c \frac{dw}{2\pi i} J^{--}(w) (G_0^+ \Phi_A(4) \tilde{G}_0^+ \Phi_B(1) + \tilde{G}_0^+ \Phi_A(4) G_0^+ \Phi_B(1)) \\ (G_0^+ \Phi_C(2) \tilde{G}_0^+ \Phi_D(3) + \tilde{G}_0^+ \Phi_C(2) G_0^+ \Phi_D(3)) \rangle_W \end{aligned}$$

where the contour c is that of Figure 2.

After performing the Giddings map of Figure 2 to the upper half plane using the ordering $[(1), (2), (3), (4)] \rightarrow [-\alpha, \alpha, \alpha^{-1}, -\alpha^{-1}]$,

$$\begin{aligned} \mathcal{A}_s = & -\frac{1}{4} \int_0^\delta d\alpha \left(\frac{d\tau}{d\alpha} \right) \left\langle \int_{\tilde{c}} \frac{dz}{2\pi i} \left(\frac{dz}{dw} \right) J^{--}(z) \right. \\ & (G_0^+ \Phi_A(-\alpha^{-1}) \tilde{G}_0^+ \Phi_B(-\alpha) + \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) G_0^+ \Phi_B(-\alpha)) \\ & \left. (G_0^+ \Phi_C(\alpha) \tilde{G}_0^+ \Phi_D(\alpha^{-1}) + \tilde{G}_0^+ \Phi_C(\alpha) G_0^+ \Phi_D(\alpha^{-1})) \right\rangle \end{aligned} \quad (5)$$

where $\tilde{c}$ is the Giddings map of the contour c and we have used that J^{--} has conformal weight 2 so $J^{--}(z) = (\frac{dz}{dw})^2 J^{--}(w)$. The overall minus sign comes from the fact that α decreases as τ increases.

4.2 Computation of t-channel contribution

Performing a re-identification of the external strings with the external legs, one finds that the t-channel contribution to the scattering amplitude is

$$\begin{aligned} \mathcal{A}_t = & \frac{1}{4} \int_0^\infty d\tau \left\langle \int_c \frac{dw}{2\pi i} J^{--}(w) \right. \\ & (G_0^+ \Phi_D(4) \tilde{G}_0^+ \Phi_A(1) + \tilde{G}_0^+ \Phi_D(4) G_0^+ \Phi_A(1)) \\ & (G_0^+ \Phi_B(2) \tilde{G}_0^+ \Phi_C(3) + \tilde{G}_0^+ \Phi_B(2) G_0^+ \Phi_C(3)) \left. \right\rangle_W \\ = & \frac{1}{4} \int_\delta^1 d\alpha \left(\frac{d\tau}{d\alpha} \right) \left\langle \int_{\tilde{c}} \frac{dz}{2\pi i} \left(\frac{dz}{dw} \right) J^{--}(z) \right. \\ & (G_0^+ \Phi_D(\alpha^{-1}) \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) + \tilde{G}_0^+ \Phi_D(\alpha^{-1}) G_0^+ \Phi_A(-\alpha^{-1})) \\ & \left. (G_0^+ \Phi_B(-\alpha) \tilde{G}_0^+ \Phi_C(\alpha) + \tilde{G}_0^+ \Phi_B(-\alpha) G_0^+ \Phi_C(\alpha)) \right\rangle \end{aligned} \quad (6)$$

where we have now used the Giddings map with the ordering $[(1), (2), (3), (4)] \rightarrow [-\alpha^{-1}, -\alpha, \alpha, \alpha^{-1}]$,

If there were no G_0^+ and $\tilde{G}_0^+$ operators, one could sum $\mathcal{A}_s$ and $\mathcal{A}_t$ to get an integral $\int_0^1 d\alpha f(\alpha)$. This is what happens in the open bosonic string four-point amplitude where $\int_0^1 d\alpha f(\alpha)$ can be related to the Veneziano amplitude using the fact that

$$\left(\frac{d\tau}{d\alpha} \right) \int_{\tilde{c}} \frac{dz}{2\pi i} \left(\frac{dz}{dw} \right) b(z) = \int d^2 z \mu_\alpha(z, \bar{z}) b(z)$$

where μ_α is the Beltrami differential corresponding to the α modulus [11]. However, for the open superstring four-point amplitude, one first has to perform contour deformations of the G_0^+ and $\tilde{G}_0^+$ operators in $\mathcal{A}_t$ until they appear in the same manner as in the expression for $\mathcal{A}_s$. As will now be shown, these contour deformations produce a *finite* contact term which is cancelled by the contribution from the quartic vertex.

Consider the expression

$$Y = \frac{d\tau}{d\alpha} \langle \int_{\tilde{c}} \frac{dz}{2\pi i} \left(\frac{dz}{dw} \right) J^{--}(z) G_0^+ \Phi_D(\alpha^{-1}) \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) \tilde{G}_0^+ \Phi_B(-\alpha) G_0^+ \Phi_C(\alpha) \rangle$$

$$= - \langle \left(\int d^2 z \mu_\alpha(z, \bar{z}) J^{--}(z) \right) \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) \tilde{G}_0^+ \Phi_B(-\alpha) G_0^+ \Phi_C(\alpha) G_0^+ \Phi_D(\alpha^{-1}) \rangle$$

which is one of the terms in $\mathcal{A}_t$. Deforming the G^+ contour off of $\Phi_D(\alpha^{-1})$, one gets

$$Y = - \langle \left(\int d^2 z \mu_\alpha(z, \bar{z}) \tilde{G}^-(z) \right) \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) \tilde{G}_0^+ \Phi_B(-\alpha) G_0^+ \Phi_C(\alpha) \Phi_D(\alpha^{-1}) \rangle.$$

Now deform the contour of $\tilde{G}^+$ off of $\Phi_B(-\alpha)$ to get

$$Y = - \langle \left(\int d^2 z \mu_\alpha(z, \bar{z}) \tilde{G}^-(z) \right) \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) \Phi_B(-\alpha) G_0^+ \Phi_C(\alpha) \tilde{G}_0^+ \Phi_D(\alpha^{-1}) \rangle$$

$$+ \langle \left(\int d^2 z \mu_\alpha(z, \bar{z}) T(z) \right) \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) \Phi_B(-\alpha) G_0^+ \Phi_C(\alpha) \Phi_D(\alpha^{-1}) \rangle$$

where T is the stress-tensor. Finally, writing $\tilde{G}^-(z) = [G_0^+, J^{--}(z)]$ and deforming the G^+ contour off of $J^{--}(z)$, one gets

$$Y = - \langle \left(\int d^2 z \mu_\alpha(z, \bar{z}) J^{--}(z) \right) \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) G_0^+ \Phi_B(-\alpha) G_0^+ \Phi_C(\alpha) \tilde{G}_0^+ \Phi_D(\alpha^{-1}) \rangle \quad (7)$$

$$+ \frac{\partial}{\partial \alpha} \langle \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) \Phi_B(-\alpha) G_0^+ \Phi_C(\alpha) \Phi_D(\alpha^{-1}) \rangle$$

where we have used that $\int d^2 z \mu_\alpha(z, \bar{z}) T(z)$ has the effect of taking the variation with respect to the modulus α .

Similarly,

$$\frac{d\tau}{d\alpha} \langle \int_{\tilde{c}} \frac{dz}{2\pi i} \left(\frac{dz}{dw} \right) J^{--}(z) \tilde{G}_0^+ \Phi_D(\alpha^{-1}) G_0^+ \Phi_A(-\alpha^{-1}) G_0^+ \Phi_B(-\alpha) \tilde{G}_0^+ \Phi_C(\alpha) \rangle \quad (8)$$

$$\begin{aligned}
&= -\langle \left(\int d^2 z \mu_\alpha(z, \bar{z}) J^{--}(z) \right) \\
&\quad G_0^+ \Phi_A(-\alpha^{-1}) \tilde{G}_0^+ \Phi_B(-\alpha) \tilde{G}_0^+ \Phi_C(\alpha) G_0^+ \Phi_D(\alpha^{-1}) \rangle \\
&\quad - \frac{\partial}{\partial \alpha} \langle G_0^+ \Phi_A(-\alpha^{-1}) \Phi_B(-\alpha) \tilde{G}_0^+ \Phi_C(\alpha) \Phi_D(\alpha^{-1}) \rangle.
\end{aligned}$$

Plugging (7) and (8) into (6), one finds

$$\begin{aligned}
\mathcal{A}_t &= -\frac{1}{4} \int_\delta^1 d\alpha \langle \left(\int d^2 z \mu_\alpha(z, \bar{z}) J^{--}(z) \right) \\
&\quad (G_0^+ \Phi_A(-\alpha^{-1}) \tilde{G}_0^+ \Phi_B(-\alpha) + \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) G_0^+ \Phi_B(-\alpha)) \\
&\quad (G_0^+ \Phi_C(\alpha) \tilde{G}_0^+ \Phi_D(\alpha^{-1}) + \tilde{G}_0^+ \Phi_C(\alpha) G_0^+ \Phi_D(\alpha^{-1})) \rangle \\
&\quad + \frac{1}{4} \int_\delta^1 d\alpha \frac{\partial}{\partial \alpha} \langle \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) \Phi_B(-\alpha) G_0^+ \Phi_C(\alpha) \Phi_D(\alpha^{-1}) \\
&\quad - G_0^+ \Phi_A(-\alpha^{-1}) \Phi_B(-\alpha) \tilde{G}_0^+ \Phi_C(\alpha) \Phi_D(\alpha^{-1}) \rangle.
\end{aligned} \tag{9}$$

Comparing with (5), one finds

$$\begin{aligned}
\mathcal{A}_s + \mathcal{A}_t &= -\frac{1}{4} \int_0^1 d\alpha \langle \left(\int d^2 z \mu_\alpha(z, \bar{z}) J^{--}(z) \right) \\
&\quad (G_0^+ \Phi_A(-\alpha^{-1}) \tilde{G}_0^+ \Phi_B(-\alpha) + \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) G_0^+ \Phi_B(-\alpha)) \\
&\quad (G_0^+ \Phi_C(\alpha) \tilde{G}_0^+ \Phi_D(\alpha^{-1}) + \tilde{G}_0^+ \Phi_C(\alpha) G_0^+ \Phi_D(\alpha^{-1})) \rangle \\
&\quad - \frac{1}{4} \langle \tilde{G}_0^+ \Phi_A(-\delta^{-1}) \Phi_B(-\delta) G_0^+ \Phi_C(\delta) \Phi_D(\delta^{-1}) \\
&\quad - G_0^+ \Phi_A(-\delta^{-1}) \Phi_B(-\delta) \tilde{G}_0^+ \Phi_C(\delta) \Phi_D(\delta^{-1}) \rangle.
\end{aligned} \tag{10}$$

Note that we have used the “cancelled propagator” argument of [13] to ignore the surface term coming from $\alpha = 1$, but this argument cannot be used to ignore the surface term coming from $\alpha = \delta$.

4.3 Computation of quartic vertex contribution

It will now be shown that this surface term from $\alpha = \delta$ is precisely cancelled by the contribution $\mathcal{A}_q$ from the quartic vertex. Using the action of (3) and including only contributions with the cyclic order $[A, B, C, D]$,

$$\mathcal{A}_q = - \sum_{cyclic} \frac{1}{24} \langle (\Phi_A(-\delta^{-1}) G_0^+ \Phi_B(-\delta) - G_0^+ \Phi_A(-\delta^{-1}) \Phi_B(-\delta)) \quad (11)$$

$$(\Phi_C(\delta) \tilde{G}_0^+ \Phi_D(\delta^{-1}) - \tilde{G}_0^+ \Phi_C(\delta) \Phi_D(\delta^{-1})) \rangle$$

where $\sum_{cyclic}$ denotes a sum over the four cyclic permutations of $[A, B, C, D]$. Using properties of the cyclic sum, the sixteen terms in $\mathcal{A}_q$ can be reordered as

$$\mathcal{A}_q = - \sum_{cyclic} \frac{1}{24} \langle 2\Phi_A G_0^+ \Phi_B \Phi_C \tilde{G}_0^+ \Phi_D$$

$$- \Phi_A G_0^+ \Phi_B \tilde{G}_0^+ \Phi_C \Phi_D - G_0^+ \Phi_A \Phi_B \Phi_C \tilde{G}_0^+ \Phi_D \rangle$$

where we have suppressed the z location of the four external strings. Note that the quartic vertex is invariant under cyclic rotations of these z locations. Deforming the contour of $\tilde{G}^+$ off of Φ_C in the second term, one gets

$$\mathcal{A}_q = - \sum_{cyclic} \frac{1}{24} \langle 3\Phi_A G_0^+ \Phi_B \Phi_C \tilde{G}_0^+ \Phi_D \quad (12)$$

$$- \tilde{G}_0^+ \Phi_A G_0^+ \Phi_B \Phi_C \Phi_D - G_0^+ \Phi_A \Phi_B \Phi_C \tilde{G}_0^+ \Phi_D \rangle.$$

But the last two terms of (12) cancel each other after summing over cyclic permutations, so

$$\mathcal{A}_q = - \sum_{cyclic} \frac{1}{8} \langle \Phi_A G_0^+ \Phi_B \Phi_C \tilde{G}_0^+ \Phi_D \rangle \quad (13)$$

$$= -\frac{1}{8} \langle \Phi_A G_0^+ \Phi_B \Phi_C \tilde{G}_0^+ \Phi_D - \Phi_A \tilde{G}_0^+ \Phi_B \Phi_C G_0^+ \Phi_D \rangle + \frac{X}{8}$$

where

$$X = \langle \tilde{G}_0^+ \Phi_A \Phi_B G_0^+ \Phi_C \Phi_D - G_0^+ \Phi_A \Phi_B \tilde{G}_0^+ \Phi_C \Phi_D \rangle.$$

Deforming the G^+ contour off of Φ_B in the first term and then deforming the $\tilde{G}^+$ contour off of Φ_D , one finds

$$\begin{aligned}
\mathcal{A}_q &= -\frac{1}{8} \langle -G_0^+ \Phi_A \Phi_B \Phi_C \tilde{G}_0^+ \Phi_D \\
&\quad - \Phi_A \Phi_B G_0^+ \Phi_C \tilde{G}_0^+ \Phi_D - \Phi_A \tilde{G}_0^+ \Phi_B \Phi_C G_0^+ \Phi_D \rangle + \frac{X}{8} \\
&= -\frac{1}{8} \langle G_0^+ \Phi_A \tilde{G}_0^+ \Phi_B \Phi_C \Phi_D - \Phi_A \tilde{G}_0^+ \Phi_B G_0^+ \Phi_C \Phi_D \\
&\quad - \Phi_A \tilde{G}_0^+ \Phi_B \Phi_C G_0^+ \Phi_D \rangle + \frac{X}{4}.
\end{aligned}$$

Finally, deforming the G_0^+ contour off of Φ_D in the last term, one finds

$$\mathcal{A}_q = \frac{X}{4}. \quad (14)$$

So combining (14) with (10),

$$\begin{aligned}
\mathcal{A} &= \mathcal{A}_s + \mathcal{A}_t + \mathcal{A}_q = -\frac{1}{4} \int_0^1 d\alpha \langle \left(\int d^2 z \mu_\alpha(z, \bar{z}) J^{--}(z) \right) \\
&\quad (G_0^+ \Phi_A(-\alpha^{-1}) \tilde{G}_0^+ \Phi_B(-\alpha) + \tilde{G}_0^+ \Phi_A(-\alpha^{-1}) G_0^+ \Phi_B(-\alpha)) \\
&\quad (G_0^+ \Phi_C(\alpha) \tilde{G}_0^+ \Phi_D(\alpha^{-1}) + \tilde{G}_0^+ \Phi_C(\alpha) G_0^+ \Phi_D(\alpha^{-1})) \rangle \\
&= \int_0^1 d\alpha \langle \left(\int d^2 z \mu_\alpha(z, \bar{z}) G^-(z) \right) \\
&\quad \Phi_A(-\alpha^{-1}) G_0^+ \Phi_B(-\alpha) G_0^+ \Phi_C(\alpha) \tilde{G}_0^+ \Phi_D(\alpha^{-1}) \rangle
\end{aligned} \quad (15)$$

where we have used contour deformations of G^+ and $\tilde{G}^+$ to write all four terms in the same form. Note that there are no surface term contributions coming from these deformations due to the “cancelled propagator” argument [13] which can be used when $\alpha = 0$ or $\alpha = 1$.

So we have proven our claim that, after including the contribution from the quartic vertex of the action of (1), the second-quantized prescription agrees with the first-quantized prescription for on-shell tree-level four-point open superstring scattering amplitudes.

Acknowledgements: The authors would like to thank CNPq grant 300256/94-9 and FAPESP grant 96/12280-0 for financial support.

References

- [1] C. Wendt, Nucl. Phys. **B314** (1989) 209.
- [2] E. Witten, Nucl. Phys. **B276** (1986) 291.
- [3] C.R. Preitschopf, C.B. Thorn, and S. Yost, Nucl. Phys. **B337** (1990) 363; I. Ya. Aref'eva, P.B. Medvedev, and A.P. Zubarev, Nucl. Phys. **B341** (1990) 464; I. Ya. Aref'eva and P.B. Medvedev, Phys. Lett. **B202** (1988) 510; N. Berkovits, M. Hatsuda and W. Siegel, Nucl. Phys. **B371** (1992) 434, hep-th/9108021.
- [4] J. Greensite and F.R. Klinkhamer, Nucl. Phys. **B291** (1987) 557; J. Greensite and F. Klinkhamer, Nucl. Phys. **B304** (1988) 108.
- [5] N. Berkovits, Nucl. Phys. **B450** (1995) 90.
- [6] E. Witten, Nucl. Phys. **B268** (1986) 253.
- [7] N. Berkovits and C. Vafa, Nucl. Phys. **B433** (1995) 123, hep-th 9407190.
- [8] S. Donaldson, Proc. Lond. Math. Soc. **50** (1985) 1; V.P. Nair and J. Schiff, Phys. Lett. **B246** (1990) 423.
- [9] N. Berkovits, Nucl. Phys. **B431** (1994) 258, hep-th 9404162.
- [10] N. Berkovits and M.M. Leite, Phys. Lett. **B454** (1999) 38, hep-th 9812153.
- [11] S. Giddings, Nucl. Phys. **B278** (1986) 242.
- [12] D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. **B271** (1986) 93.
- [13] J. Polchinski, **String Theory, vol.1**, Cambridge University Press (1998).